
Lattice Cryptography (2. Encryption Using Polynomial Rings)


Vadim Lyubashevsky

IBM Research Europe, Zurich

Encryption Scheme



1. Multiplication As takes quadratic time (not so fast in practice)



2. <r,t> is an integer, and so we can only encrypt 1 bit at a time

Use Polynomials

```
f(x) is a polynomial x^n + a_{n-1}x^{n-1} + ... + a_1x + a_0
```

 $R = Z_p[x]/(f(x))$ is a polynomial ring with

- Addition mod p
- Polynomial multiplication mod p and f(x)

Each element of R consists of n elements in Z_p

In R:

- small+small = small
- small*small = small (depending on f(x))

Example ring $Z_{17}[X]/(X^4+1)$

Elements are
$$z(X)=z_3X^3+z_2X^2+z_1X+z_0$$

where z_i are integers mod 17

Addition is the usual coordinate-wise addition

Multiplication is the usual polynomial multiplication followed by reduction modulo X^4+1


Multiplication and coefficient growth in $Z[X]/(X^4+1)$

Polynomial multiplication as Matrix-Vector multiplication


$$a(X)=a_3X^3+a_2X^2+a_1X+a_0$$

 $b(x)=b_3X^3+b_2X^2+b_1X+b_0$

$$a(X)*b(X) = b(X)*a_0 + (b(X) * X) * a_1 + (b(X)*X^2) * a_2 + (b(X)*X^3) * a_3$$



when the ring is $Z[X]/(X^4+1)$



 a_3


because $b(X)^*X^i$ has small coefficients, the coefficients grow "slowly" during multiplication if the ring were $Z[X]/(X^n + 2X^{n-1} + 1)$, then $b(X)^*X^{n-1}$ can have coefficients 2^n times larger than b(X)

Operations

Basic Computational Domain:


Polynomial ring
$$Z_p[X]/(X^n+1)$$

Operations used in the schemes: • and * in the ring:




Hard problem LWE over Rings

Basic Hard Problem:




Encryption Scheme



Encryption scheme over $Z_q[X]/(X^n+1)$


Key Generation



Decryption

Round
$$\begin{bmatrix} v & - & u_1 & u_2 & s_1 \\ & s_2 & & & \end{bmatrix} = \begin{bmatrix} m & s_1 & & \\ & s_2 & & & \end{bmatrix}$$

Encryption



Operations in $R=Z_p[X]/(X^n+1)$

- Additions are easy and cheap
- To get cheap multiplications, we pick p = 1 (mod 2n)

Can then write
$$X^n+1 = (X-r_1)(X-r_2) \cdot \cdot \cdot (X-r_n) \mod p$$

Polynomials in R can be represented in two ways

Coefficient representation

$$f = a_0 + a_1 x + \dots + a_{n-1} x^{n-1}$$

Chinese Remainder Theorem (CRT) representation

$$\hat{f} = (f(r_1), f(r_2), ..., f(r_n))$$

Multiplication using NTT (Number Theoretic Transform)

Want to multiply f * g, which are in coefficient representation

1.
$$f \to \hat{f} = (f(r_1), f(r_2), ..., f(r_n))$$

2. $g \to \hat{g} = (g(r_1), g(r_2), ..., g(r_n))$
3. $\widehat{f * g} = (f(r_1) * g(r_1), ..., f(r_n) * g(r_n))$
4. $\widehat{f * g} \to f * g$

Example in $R=Z_{17}[x]/(X^4+1)$


$$x^4 + 1 = (x - 2)(x + 2)(x - 8)(x + 8) \bmod 17$$

We will show how to efficiently convert between the coefficient and CRT representations

This is the main part of polynomial multiplication

$$f \rightarrow \hat{f}$$

$$f \bmod (x^4 + 1)$$




$$f_{11} = f \ mod \ (x^2 - 4)$$

$$f_{12} = f \ mod \ (x^2 + 4)$$



 $f_{21} = f_{11} \bmod (x - 2)$









$$f_{23} = f_{12} \ mod \ (x - 8)$$

$$f_{24} = f_{12} \bmod (x+8)$$

$$f(-2)$$

$$f \rightarrow \hat{f}$$

$$f \bmod (x^4 + 1)$$

$$1-4x+3x^2-5x^3$$

$$f_{11} = f \ mod \ (x^2 - 4)$$

$$1 - 4x + 3 * 4 - 5 * 4x$$



$$f_{21} = f_{11} \bmod (x-2)$$

$$|f_{21} = f_{11} \mod (x-2)|$$
 $|f_{22} = f_{11} \mod (x+2)|$

$$f_{23} = f_{12} \bmod$$

$$f_{23} = f_{12} \ mod \ (x - 8)$$

 $f_{12} = f \ mod \ (x^2 + 4)$

1 - 4x - 3 * 4 + 5 * 4x

$$f_{24} = f_{12} \bmod (x+8)$$

$$f(-2)$$

$$f(-8)$$


$$f \rightarrow \hat{f}$$

$$f \bmod (x^4 + 1)$$

$$1-4x+3x^2-5x^3$$

$$f_{11} = f \ mod \ (x^2 - 4)$$

$$-4 - 7x$$



$$6-x$$







$$f_{21} = f_{11} \bmod (x-2)$$

$$f_{22} = f_{11} \bmod (x+2)$$

$$f_{23} = f_{12} \ mod \ (x - 8)$$

$$f_{24} = f_{12} \bmod (x+8)$$

$$-2$$

$$-3$$

$$f(-2)$$

$$f(-8)$$

$$\hat{f} \rightarrow f$$

$$f \bmod (x^4 + 1)$$

$$f_{11} = f \ mod \ (x^2 - 4)$$

$$f_{12} = f \bmod (x^2 + 4)$$

$$a + bx$$

$$a + bx \qquad a + 2b = -1$$

$$a-2b=-7$$

$$|f_{21} = f_{11} \bmod (x-2)|$$

$$f_{22} = f_{11} \bmod (x+2)$$

$$f_{23} = f_{12} \bmod (x - 8)$$

$$f_{24} = f_{12} \bmod (x+8)$$

f(2)

f(8)


$$\hat{f} \rightarrow f$$

$$f \bmod (x^4 + 1)$$

$$1-4x+3x^2-5x^3$$


$$|f_{11} = f \bmod (x^2 - 4)|$$


$$-4 - 7x$$



$$6-x$$







$$f_{24} = f_{12} \bmod (x+8)$$

_1

 $f_{21} = f_{11} \bmod (x-2)$

—7

-2

-3

f(2)

f(-2)

f(8)

f(-8)

Time complexity

$$X^n - r \equiv (X^{n/2} - \sqrt{r})(X^{n/2} + \sqrt{r})$$

$$a = \sum_{i=0}^{n-1} a_i X^i,$$

$$a \mod X^{n/2} - \sqrt{r} = \sum_{i=0}^{n/2-1} b_i X^i,$$

$$a \mod X^{n/2} + \sqrt{r} = \sum_{i=0}^{n/2-1} c_i X^i,$$

Computing the NTT

Computing the inverse NTT

$$b_{i} = a_{i} + \sqrt{r} \cdot a_{i+n/2}$$

$$c_{i} = a_{i} - \sqrt{r} \cdot a_{i+n/2}$$

$$2 \cdot a_{i} = b_{i} + c_{i}$$

$$2 \cdot a_{i+n/2} = (\sqrt{r})^{-1} \cdot (b_{i} - c_{i})$$

n additions and n/2 multiplications

n additions and n/2 multiplications

Computation time for $R=Z_p[x]/(x^n+1)$

```
f \to \hat{f} and \hat{f} \to f log n levels (n/2 multiplications) and (n additions) modulo p per level
```

Problem session

Implement NTT and NTT⁻¹ over $Z_{257}[X]/(X^{32} + 1)$

- Create a tree (given to you) of roots and inverse roots that will be used for all NTT and NTT⁻¹ computations
- All the operations are to be done "in place" -- do not create new vectors
 - At level 0, your vector consists of a polynomial with 32 coefficients
 - At level 1, your vector consists of 2 polynomials with 16 coefficients
 - •
 - At the bottom level, your vector consists of 32 polynomials with 1 coefficient

Supplementary reading: Section 4 of https://github.com/VadimLyubash/LatticeTutorial