

# Multivariate cryptography - Cryptanalysis techniques II

SLMath summer school:

Introduction to Quantum-Safe Cryptography (IBM Zurich)

Simona Samardjiska

July, 2024

Institute for Computing and Information Sciences Radboud University

#### Recall the MinRank problem

MinRank  $MR(n, m, r, M_1, \ldots, M_m)$ 

**Input**:  $n, m, r \in \mathbb{N}$ , and  $M_1, \ldots, M_m \in \mathcal{M}_n(\mathbb{F}_q)$ .

**Question**: Find – if any – a nonzero *m*-tuple  $(\lambda_1, \ldots, \lambda_m) \in \mathbb{F}_q^m$  s.t.:


$$\mathsf{Rank}\left(\sum_{i=1}^m \lambda_i \, M_i\right) \leqslant r.$$

[Courtois '01], [Buss & Shallit '99]


$$\mathcal{P} = (p_1, p_2, \dots, p_m)$$
 - public polynomials,  
 $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .




$$\mathcal{P}=(p_1,p_2,\ldots,p_m)$$
 - public polynomials,  $\mathbf{P}_1,\mathbf{P}_2,\ldots,\mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .




$$\mathcal{P} = (p_1, p_2, \dots, p_m)$$
 - public polynomials,  $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .




$$\mathcal{P} = (p_1, p_2, \dots, p_m)$$
 - public polynomials,  $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .



$$\mathcal{P} = (p_1, p_2, \dots, p_m)$$
 - public polynomials,  $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .



$$\mathcal{P}=(p_1,p_2,\ldots,p_m)$$
 - public polynomials,  $\mathbf{P}_1,\mathbf{P}_2,\ldots,\mathbf{P}_m$  - matrix representations of the coordinates of  $\mathcal{P}$ .



### Baby example UOV

Similar approach works for UOV, although it is not a result of "rank defect" (at leaset not so obvious)

$$f_1(x_1, \dots, x_6) = x_1 x_2 + x_2 x_4 + x_3 x_6 + x_4 x_6 + x_5 x_6 + x_6$$

$$f_2(x_1, \dots, x_6) = x_1 x_4 + x_3 x_4 + x_3 x_6 + x_4 x_6 + x_6$$

$$f_3(x_1, \dots, x_6) = x_2 x_3 + x_3 x_5 + x_2 x_4 + x_2 x_6 + x_4 x_5 + x_1 x_6 + x_4 x_6 + x_5 x_6$$

### Baby example UOV

Similar approach works for UOV, although it is not a result of "rank defect" (at leaset not so obvious)

$$f_1(x_1, \dots, x_6) = x_1x_2 + x_2x_4 + x_3x_6 + x_4x_6 + x_5x_6 + x_6$$

$$f_2(x_1, \dots, x_6) = x_1x_4 + x_3x_4 + x_3x_6 + x_4x_6 + x_6$$

$$f_3(x_1, \dots, x_6) = x_2x_3 + x_3x_5 + x_2x_4 + x_2x_6 + x_4x_5 + x_1x_6 + x_4x_6 + x_5x_6$$

$$\overline{S}'$$
:  $x_4 \rightarrow x_4 + x_6$   
 $x_2 \rightarrow x_2 + x_5$ 

### Baby example UOV

Similar approach works for UOV, although it is not a result of "rank defect" (at leaset not so obvious)

$$f_1(x_1, \dots, x_6) = x_1x_2 + x_2x_4 + x_3x_6 + x_4x_6 + x_5x_6 + x_6$$

$$f_2(x_1, \dots, x_6) = x_1x_4 + x_3x_4 + x_3x_6 + x_4x_6 + x_6$$

$$f_3(x_1, \dots, x_6) = x_2x_3 + x_3x_5 + x_2x_4 + x_2x_6 + x_4x_5 + x_1x_6 + x_4x_6 + x_5x_6$$

$$\overline{S}'$$
:  $x_4 \rightarrow x_4 + x_6$   
 $x_2 \rightarrow x_2 + x_5$ 

After change of variables, we have separated (some) of the oil space( $x_5, x_6$ ):

$$f_1(x_1, \dots, x_6) = x_1 x_2 + x_1 x_5 + x_2 x_4 + x_2 x_6 + x_4 x_5 + x_3 x_6 + x_4 x_6$$


$$f_2(x_1, \dots, x_6) = x_1 x_4 + x_1 x_6 + x_3 x_4 + x_4 x_6$$


$$f_3(x_1, \dots, x_6) = x_2 x_3 + x_2 x_4 + x_4 x_6 + x_1 x_6 + x_6$$

### **UOV** partial revealing of structure ("good keys")

UOV

$$f_s(x) = \sum_{i \in V, j \in V} \gamma_{ij}^{(s)} x_i x_j + \sum_{i \in V, j \in O} \gamma_{ij}^{(s)} x_i x_j,$$







Good Keys for UOV

## Rainbow partial revealing of structure ("good keys")

Rainbow before and after applying an input and output change of basis (separating a a good key)





Good key for Rainbow -

# Measuring linear spaces

## **Linear spaces of** (n, m)-functions

- **Differential** of f:  $\mathcal{D}_w f(x) = f(x+w) f(x) f(w) + f(0)$
- Linearity for (n, m) functions  $f : \mathbb{F}_q^n \to \mathbb{F}_q^m$  defined already 1992 by Nyberg
- $w \in \mathbb{F}_q^n$  linear structure of f if

$$\mathcal{D}_w f(x) = 0 \quad \forall \ x \in \mathbb{F}_q^n$$

• Linear space of f - generated by the linear structures of f.

Quadratic form  $f: \mathcal{D}_w f(x) = w^T F x$ , for a symmetric matrix F,

• Ker(**F**) - **linear space of** *f* .

[Nyberg92] Quadratic (n, m)-function f

• Linearity - measured using the smallest rank r of any of the components  $w^{T} \cdot f$ .

#### Maximum nonlinearity

- Bent functions Rank $(F_w) = n$ , even  $n, m \le n/2$ ,
- Almost bent (AB) functions Rank $(F_w) = n 1$ , odd n = m

## **Linear spaces of** (n, m)-functions

- **Differential** of  $f: \mathcal{D}_w f(x) = f(x+w) f(x) f(w) + f(0)$
- ullet Linearity for (n,m) functions  $f:\mathbb{F}_q^n o \mathbb{F}_q^m$  defined already 1992 by Nyberg
- $w \in \mathbb{F}_q^n$  linear structure of f if

$$\mathcal{D}_w f(x) = 0 \quad \forall \ x \in \mathbb{F}_q^n$$
.

• **Linear space** of f - generated by the linear structures of f.

Quadratic form  $f: \mathcal{D}_w f(x) = w^{\mathsf{T}} \mathbf{F} x$ , for a symmetric matrix  $\mathbf{F}$ ,

• Ker(F) - linear space of f.

[Nyberg92] **Quadratic** (n, m)-function f:

• Linearity - measured using the smallest rank r of any of the components  $w^{\mathsf{T}} \cdot f$ .

#### Maximum nonlinearity

- Bent functions Rank( $F_w$ ) = n, even n,  $m \le n/2$ ,
- Almost bent (AB) functions Rank( $F_w$ ) = n-1, odd n=m.

## **Linear spaces of** (n, m)-functions

- **Differential** of f:  $\mathcal{D}_w f(x) = f(x+w) f(x) f(w) + f(0)$
- Linearity for (n, m) functions  $f : \mathbb{F}_q^n \to \mathbb{F}_q^m$  defined already 1992 by Nyberg
- $w \in \mathbb{F}_q^n$  linear structure of f if

$$\mathcal{D}_w f(x) = 0 \quad \forall \ x \in \mathbb{F}_q^n$$
.

• **Linear space** of f - generated by the linear structures of f.

Quadratic form  $f: \mathcal{D}_w f(x) = w^{\mathsf{T}} \mathbf{F} x$ , for a symmetric matrix  $\mathbf{F}$ ,

• Ker(F) - linear space of f.

[Nyberg92] **Quadratic** (n, m)-function f:

• Linearity - measured using the smallest rank r of any of the components  $w^{\mathsf{T}} \cdot f$ .

#### Maximum nonlinearity:

- Bent functions Rank $(\mathbf{F}_w) = n$ , even  $n, m \leq n/2$ ,
- Almost bent (AB) functions Rank( $\mathbf{F}_w$ ) = n-1, odd n=m.

#### Example 1:

#### Example 1:

 $(1,0,0,1)^{\mathsf{T}}$ 

$$f: f_1 = x_1x_2 + x_3 f_2 = x_1x_3 + x_2 + x_3 f_3 = x_2x_3 + x_1 + x_2 + x_3 f_4 = x_1x_2$$

 $\cdot f$  is linear

$$f':$$

$$f'_{1} = x_{1}x_{2} + x_{3}$$

$$f'_{2} = x_{1}x_{2} + x_{2} + x_{3}$$

$$f'_{3} = x_{2}x_{3} + x_{1} + x_{2} + x_{3}$$

$$f'_{4} = x_{1}x_{2} + x_{2}x_{3}$$

$$(1,0,1,1)^{\mathsf{T}}$$
  $f'$  is linear  $(1,1,0,0)^{\mathsf{T}}$   $f'$  is linear

Both have maximum linearity, but f' is linear on a larger space! It is an important measure!

## Oil & Vinegar maps

f :

$$f_1(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_4 + x_1x_2 + x_3$$
  
$$f_2(x_1, x_2, x_3, x_4) = x_2x_3 + x_1x_4 + x_2x_4 + x_3$$

f is linear on the oil subspace (when you fix the vinegar variables)!

$$f_1(c_1, c_2, x_3, x_4) = c_1 x_3 + c_2 x_4 + c_1 c_2 + x_3$$

$$f_2(c_1, c_2, x_3, x_4) = c_2 x_3 + c_1 x_4 + c_2 x_4 + x_3$$

## Oil & Vinegar maps

f :

$$f_1(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_4 + x_1x_2 + x_3$$
  
$$f_2(x_1, x_2, x_3, x_4) = x_2x_3 + x_1x_4 + x_2x_4 + x_3$$

f is linear on the oil subspace (when you fix the vinegar variables)!

$$f_1(c_1, c_2, x_3, x_4) = c_1 x_3 + c_2 x_4 + c_1 c_2 + x_3$$

$$f_2(c_1, c_2, x_3, x_4) = c_2 x_3 + c_1 x_4 + c_2 x_4 + x_3$$

## Oil & Vinegar maps

f :

$$f_1(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_4 + x_1x_2 + x_3$$
  
$$f_2(x_1, x_2, x_3, x_4) = x_2x_3 + x_1x_4 + x_2x_4 + x_3$$

f is linear on the oil subspace (when you fix the vinegar variables)!

$$f_1(c_1, c_2, x_3, x_4) = c_1 x_3 + c_2 x_4 + c_1 c_2 + x_3$$
  
$$f_2(c_1, c_2, x_3, x_4) = c_2 x_3 + c_1 x_4 + c_2 x_4 + x_3$$

## (s, t)-linearity of quadratic (n, m) function f

#### Boura and Canteaut FSE13:

```
f is said to be (s,t)-linear if there exist linear subspaces V \subset \mathbb{F}_q^n with \mathsf{Dim}(V) = s, W \subset \mathbb{F}_q^m with \mathsf{Dim}(W) = t, s.t.
```

 $\forall w \in W, w^{\mathsf{T}} \cdot f$  is linear on all cosets of V.

## (s,t)-linearity of quadratic (n,m) function f

#### Boura and Canteaut FSE13:

f is said to be (s,t)-linear if there exist linear subspaces  $V \subset \mathbb{F}_q^n$  with  $\mathsf{Dim}(V) = s$ ,  $W \subset \mathbb{F}_q^m$  with  $\mathsf{Dim}(W) = t$ , s.t.

 $\forall w \in W, w^{\mathsf{T}} \cdot f$  is linear on all cosets of V.

•  $f_W$  corresponding to all  $w^\intercal \cdot f$ ,  $w \in W$  can be written as

$$f_W(x,y) = M(x) \cdot y + G(x)$$

where  $\mathbb{F}_q^n = U \oplus V$ ,  $G: U \to \mathbb{F}_q^t$  and M(x) is a  $t \times s$  matrix with rows - components of linear functions over U.

### (s, t)-linearity of quadratic (n, m) function f

#### Boura and Canteaut FSE13:

f is said to be (s,t)-linear if there exist linear subspaces  $V \subset \mathbb{F}_q^n$  with  $\mathsf{Dim}(V) = s$ ,  $W \subset \mathbb{F}_q^m$  with  $\mathsf{Dim}(W) = t$ , s.t.

 $\forall w \in W, w^{\mathsf{T}} \cdot f$  is linear on all cosets of V.

•  $f_W$  corresponding to all  $w^T \cdot f$ ,  $w \in W$  can be written as

$$f_W(x,y) = M(x) \cdot y + G(x)$$

where  $\mathbb{F}_q^n = U \oplus V$ ,  $G: U \to \mathbb{F}_q^t$  and M(x) is a  $t \times s$  matrix with rows - components of linear functions over U.

• for  $w \in W$ ,  $\mathcal{D}_a w^{\mathsf{T}} \cdot f(b) = 0$ ,  $\forall a, b \in V$ .

### (s, t)-linearity of quadratic (n, m) function f

#### Boura and Canteaut FSE13:

f is said to be (s,t)-linear if there exist linear subspaces  $V \subset \mathbb{F}_q^n$  with  $\mathsf{Dim}(V) = s$ ,  $W \subset \mathbb{F}_q^m$  with  $\mathsf{Dim}(W) = t$ , s.t.

 $\forall w \in W, w^{\mathsf{T}} \cdot f$  is linear on all cosets of V.

•  $f_W$  corresponding to all  $w^T \cdot f$ ,  $w \in W$  can be written as

$$f_W(x,y) = M(x) \cdot y + G(x)$$

where  $\mathbb{F}_q^n = U \oplus V$ ,  $G: U \to \mathbb{F}_q^t$  and M(x) is a  $t \times s$  matrix with rows - components of linear functions over U.

- for  $w \in W$ ,  $\mathcal{D}_a w^\intercal \cdot f(b) = 0$ ,  $\forall a, b \in V$ .
- for  $w \in W$ ,  $f_W(0,y) = M(0) \cdot y + G(0) = 0, \forall (0,a) \in V$  all components in W vanish on the V space

#### **Example:**

$$f_1(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_4 + x_1x_2 + x_3$$
  

$$f_2(x_1, x_2, x_3, x_4) = x_2x_3 + x_1x_4 + x_2x_4 + x_3$$

$$f$$
 is  $(2,2)$ -linear,  $V = \langle (0,0,1,0), (0,0,0,1) \rangle$ ,  $W = \langle (1,0), (0,1) \rangle$ 

$$f_1(x_1, x_2, x_3, x_4) = x_1 x_3 + x_1 x_4 + x_2$$

$$f_2(x_1, x_2, x_3, x_4) = x_1 x_2 + x_1 x_4 + x_1 x_3$$

$$f_3(x_1, x_2, x_3, x_4) = x_1 x_3 + x_2 x_3 + x_2 x_4$$

$$f \text{ is } (3,2) - \text{linear,} \\ V = \langle (0,1,0,0), (0,0,1,0), (0,0,0,1) \rangle, \ W = \langle (1,0,0), (0,1,0) \rangle$$

#### **Example:**

$$f_1(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_4 + x_1x_2 + x_3$$
  
$$f_2(x_1, x_2, x_3, x_4) = x_2x_3 + x_1x_4 + x_2x_4 + x_3$$

$$f \text{ is } (2,2) - \text{linear,} \\ V = \langle (0,0,1,0), (0,0,0,1) \rangle, \ W = \langle (1,0), (0,1) \rangle \\ \\ f_1(x_1,x_2,x_3,x_4) = x_1x_3 + x_1x_4 + x_2 \\ \\ f_2(x_1,x_2,x_3,x_4) = x_1x_2 + x_1x_4 + x_1x_3 \\ \\ f_3(x_1,x_2,x_3,x_4) = x_1x_3 + x_2x_3 + x_2x_4 \\$$

$$f \text{ is } (3,2) - \text{linear,} \\ V = \langle (0,1,0,0), (0,0,1,0), (0,0,0,1) \rangle, \ W = \langle (1,0,0), (0,1,0) \rangle$$

#### Back to UOV - what does the above mean?

 $\mathcal{D}_a f(b) = 0$ ,  $\forall a, b$  in the oil space O.  $f(0, a) = 0, \forall a \in O$  - the oil and vinegar map vanishes on the oil space!

#### Basis for the new definition of UOV [Beullens21]

A consequence? - Reconciliation Attack [Ding et al.] In a nutshel: Recover (s, m) linearity of the public  $\mathcal{P}: P_1, \ldots, P_r$ 

Solve

$$x^{(j)}\mathbf{P}_{i}x^{(k)} = 0, i \in \{1, ..., m\}, j, k \in \{1, ..., s\}, j < k$$
  
 $x^{(k)}\widetilde{\mathbf{P}}_{i}x^{(k)} = 0, i \in \{1, ..., m\}, k \in \{1, ..., s\},$ 

in the unknown basis vectors  $x^{(j)}$  of the oil space O,

where 
$$\mathbf{P}_i := \widetilde{\mathbf{P}}_i + \widetilde{\mathbf{P}}_i^{\mathsf{T}}$$
.

#### Back to UOV - what does the above mean?

 $\mathcal{D}_a f(b) = 0$ ,  $\forall a, b$  in the oil space O.  $f(0, a) = 0, \forall a \in O$  - the oil and vinegar map vanishes on the oil space!

#### Basis for the new definition of UOV [Beullens21]

A consequence? - Reconciliation Attack [Ding et al.]

In a nutshel: Recover (s, m) linearity of the public  $\mathcal{P}: \mathbf{P}_1, \dots, \mathbf{P}_m$ 

Solve:

$$x^{(j)}\mathbf{P}_{i}x^{(k)} = 0, i \in \{1,...,m\}, j,k \in \{1,...,s\}, j < k$$
  
 $x^{(k)}\widetilde{\mathbf{P}}_{i}x^{(k)} = 0, i \in \{1,...,m\}, k \in \{1,...,s\},$ 

in the unknown basis vectors  $x^{(j)}$  of the oil space O,

where 
$$\mathbf{P}_i := \widetilde{\mathbf{P}}_i + \widetilde{\mathbf{P}}_i^{\mathsf{T}}$$
.

## Reconciliation attack [Ding et al.]

#### As given in [SG14]:

Solve the quadratic

$$x^{(j)}\mathbf{P}_{i}x^{(k)} = 0, i \in \{1,...,m\}, j,k \in \{1,...,c\}, j < k$$
  
 $x^{(k)}\widetilde{\mathbf{P}}_{i}x^{(k)} = 0, i \in \{1,...,m\}, k \in \{1,...,c\},$ 

in the unknown basis vectors  $x^{(k)}$  of the space O.

- $m\binom{c+1}{2}$  quadratic and bilinear equations  $\binom{n-m}{c}$  variables We must choose c s.t.  $m\binom{c+1}{2} \geq (n-m)c$  (typically at least 2)
- 2 Then solve the linear

$$x^{(j)}\mathbf{P}_{i}x^{(k)} = 0, i \in \{1,...,m\}, j \in \{1,...,c\}, k \in \{c+1,...,m\}, j < k$$

in the unknown basis vectors  $x^{(k)}$  of the oil space O.

[For first k, mc linear equations (n-m) variables Works if  $m(c+1) \ge n$ , otherwise plug in in step 1 and solve easier quadratic system]

## Making the reconciliation attack practical

#### Important about the attack:

- If c taken big enough in the first step, second step is always polynomial
- First step is the expensive one
- Questions:
  - Can we have a polynomial second step for smaller c?
  - Yes, only one vector seems to be enough!
  - Can we find easier (than step 1) vectors in the oil space?
  - Yes, intersection attack!

#### One oil vector breaks UOV!

- Shown in [Aulbach, Campos, Krämer, S, Stöttinger '23]
- Simpler view in [Pébereau'24]
  - Assume n < 3m
  - Assume an oil vector o is known
  - Recall that  $\mathcal{D}_{o}f(b) =$

$$\mathcal{D}_{o}f(b)=0$$
,  $\forall b$  in the oil space  $O$ .

so the oil space O lives in the kernel of the differential  $\mathcal{D}_{o}$ 

$$|\mathsf{Ker}(\mathcal{D}_{\boldsymbol{o}})| = n - m$$

• Restrict the public key to  $Ker(\mathcal{D}_o)$  using a basis matrix  $S_{Ker}$ 

$$\mathcal{P}_{|\operatorname{\mathsf{Ker}}(\mathcal{D}_{m{o}})} = \mathcal{P} \circ m{S}_{\operatorname{\mathsf{Ker}}}$$

- Obtain a (n m, m) UOV instance
  - Unknown oil space O' can be found by Kipnis-Shamir attack '98 (becomes polynomial)
  - Alternatively, use Step 2 of reconciliation attack for c=1 (becomes polynomial)
- Go back to original UOV instance
  - Basis of unknown oil space  $\mathbf{B}_O = \mathbf{S}_{\mathsf{Ker}} \cdot \mathbf{B}_{O'}$

### Finding one oil vector: enhancement of Kipnis-Shamir attack '98

- Kipnis-Shamir attack '98 Broke Oil & Vinegar by Patarin (n=2m)
- Recall that  $\mathcal{D}_o f(b) = 0$ ,  $\forall b$  in the oil space O.
- In matrix form

$$o^{(j)}\mathbf{P}_{i}o^{(k)} = 0, i \in \{1,...,m\}, j,k \in \{1,...,m\}$$
  
 $\mathbf{P}_{i} \cdot O \subset O^{\perp}$ 


- $\bullet |\mathbf{P}_i \cdot O| = m, |O^{\perp}| = n m$
- $|\mathbf{P}_i \cdot O \cap \mathbf{P}_j \cdot O| \ge |\mathbf{P}_i \cdot O| + |\mathbf{P}_j \cdot O| |O^{\perp}| = 3m n$

### Finding one oil vector: enhancement of Kipnis-Shamir attack '98

- Kipnis-Shamir attack '98 Broke Oil & Vinegar by Patarin (n = 2m)
- Recall that  $\mathcal{D}_{o}f(b) = 0$ ,  $\forall b$  in the oil space O.
- In matrix form

$$o^{(j)} \mathbf{P}_i o^{(k)} = 0, i \in \{1, ..., m\}, j, k \in \{1, ..., m\}$$
  
 $\mathbf{P}_i \cdot O \subset O^{\perp}$ 

- $|\mathbf{P}_i \cdot O| = m$ ,  $|O^{\perp}| = n m$
- $|\mathbf{P}_i \cdot O \cap \mathbf{P}_j \cdot O| \ge |\mathbf{P}_i \cdot O| + |\mathbf{P}_j \cdot O| |O^{\perp}| = 3m n$




### Finding one oil vector: enhancement of Kipnis-Shamir attack '98

- Kipnis-Shamir attack '98 Broke Oil & Vinegar by Patarin (n = 2m)
- Recall that  $\mathcal{D}_{o}f(b) = 0$ ,  $\forall b$  in the oil space O.
- In matrix form

$$o^{(j)} \mathbf{P}_i o^{(k)} = 0, i \in \{1, ..., m\}, j, k \in \{1, ..., m\}$$
  
 $\mathbf{P}_i \cdot O \subset O^{\perp}$ 

- $|\mathbf{P}_i \cdot O| = m$ ,  $|O^{\perp}| = n m$
- $|\mathbf{P}_i \cdot O \cap \mathbf{P}_j \cdot O| \ge |\mathbf{P}_i \cdot O| + |\mathbf{P}_j \cdot O| |O^{\perp}| = 3m n$



# Intersection attack [Beullens '20]

- Focus on n < 3m
- We want to find x in the intersection  $P_i \cdot O \cap P_j \cdot O$
- But then  $P_i^{-1}x \in O$  and  $P_i^{-1}x \in O$  are two oil vectors
- We can do the reconciliation attack but on steroids!
  - Fix 3m n coordinates of x and solve the quadratic system

$$(\mathbf{P}_{1}^{-1}x)^{\top}\mathbf{P}_{i}\mathbf{P}_{2}^{-1}x = 0, i \in \{1, ..., m\}$$

$$(\mathbf{P}_{1}^{-1}x)^{\top}\widetilde{\mathbf{P}}_{i}(\mathbf{P}_{1}^{-1}x) = 0, i \in \{1, ..., m\}$$

$$(\mathbf{P}_{2}^{-1}x)^{\top}\widetilde{\mathbf{P}}_{i}(\mathbf{P}_{2}^{-1}x) = 0, i \in \{1, ..., m\}$$

- 3m equations and 2n 3m variables
- We now have two oil vectors, the rest is easy!

# Intersection attack [Beullens '20]

- Focus on n < 3m
- We want to find x in the intersection  $P_i \cdot O \cap P_j \cdot O$
- But then  $P_i^{-1}x \in O$  and  $P_i^{-1}x \in O$  are two oil vectors
- We can do the reconciliation attack but on steroids!
  - Fix 3m n coordinates of x and solve the quadratic system

$$(\mathbf{P}_{1}^{-1}x)^{\top}\mathbf{P}_{i}\mathbf{P}_{2}^{-1}x = 0, i \in \{1, ..., m\}$$

$$(\mathbf{P}_{1}^{-1}x)^{\top}\widetilde{\mathbf{P}}_{i}(\mathbf{P}_{1}^{-1}x) = 0, i \in \{1, ..., m\}$$

$$(\mathbf{P}_{2}^{-1}x)^{\top}\widetilde{\mathbf{P}}_{i}(\mathbf{P}_{2}^{-1}x) = 0, i \in \{1, ..., m\}$$

- 3m equations and 2n 3m variables
- We now have two oil vectors, the rest is easy!

Better said, let's take a different perspective...


So far we considered m symmetric matrices representing our polynomials.

Like this:

### Better said, let's take a different perspective...

So far we considered m symmetric matrices representing our polynomials.


#### Like this:




### Better said, let's take a different perspective...

So far we considered m symmetric matrices representing our polynomials.

Like this:




But, this is also good...




### Better said, let's take a different perspective...

So far we considered m symmetric matrices representing our polynomials.


Like this:



But, this is also good...




And this!




### Better said, let's take a different perspective...

So far we considered m symmetric matrices representing our polynomials.


Like this:



But, this is also good...



And this!



- This is different tensor view, but the same object!
- Instead of array of two-dimensional matrices, we look at it as a three-dimensional qube!

Recall, UOV has an important hidden linear spaces (the oil space)...

But no rank defects!

Sure


Recall, UOV has an important hidden linear spaces (the oil space)...

But no rank defects!



Recall, UOV has an important hidden linear spaces (the oil space)...


### But no rank defects!






Recall, UOV has an important hidden linear spaces (the oil space)...

### But no rank defects!










Recall, UOV has an important hidden linear spaces (the oil space)...

#### But no rank defects!










- In the second and third view, we observe a rank defect!
- We can use (Rectangular) MinRank!
- Beullens '22 used it to improve the analysis on UOV and Rainbow
- **Important takeaway:** The two types of important linear spaces can be characterized in the same way!

Recall, UOV has an important hidden linear spaces (the oil space)...

#### But no rank defects!










- In the second and third view, we observe a rank defect!
- We can use (Rectangular) MinRank!
- Beullens '22 used it to improve the analysis on UOV and Rainbow
- **Important takeaway:** The two types of important linear spaces can be characterized in the same way!

Recall, UOV has an important hidden linear spaces (the oil space)...

#### But no rank defects!







- In the second and third view, we observe a rank defect!
- We can use (Rectangular) MinRank!
- Beullens '22 used it to improve the analysis on UOV and Rainbow
- Important takeaway: The two types of important linear spaces can be characterized in the same way!

- MQ-Sign submitted to the Korean PQC competition for standardization
- Now a finalist
- MQ-Sign design principle:
  - UOV map
  - Sparse polynomials to reduce key size only v coefficients per polynomial
  - Four variants with different level of sparsness
  - Equivalent keys technique to reduce key size
- Polynomial time attack [AST23] on the sparse variants
- Result of flawed use of equivalent keys

- MQ-Sign submitted to the Korean PQC competition for standardization
- Now a finalist
- MQ-Sign design principle:
  - UOV map
  - Sparse polynomials to reduce key size only v coefficients per polynomial
  - Four variants with different level of sparsness
  - Equivalent keys technique to reduce key size
- Polynomial time attack [AST23] on the sparse variants
- Result of flawed use of equivalent keys

- MQ-Sign submitted to the Korean PQC competition for standardization
- Now a finalist
- MQ-Sign design principle:
  - UOV map
  - Sparse polynomials to reduce key size only v coefficients per polynomial
  - · Four variants with different level of sparsness
  - Equivalent keys technique to reduce key size
- Polynomial time attack [AST23] on the sparse variants
- Result of flawed use of equivalent keys

- MQ-Sign submitted to the Korean PQC competition for standardization
- Now a finalist
- MQ-Sign design principle:
  - UOV map
  - Sparse polynomials to reduce key size only v coefficients per polynomial
  - · Four variants with different level of sparsness
  - Equivalent keys technique to reduce key size
- Polynomial time attack [AST23] on the sparse variants
- Result of flawed use of equivalent keys

## **MQ-Sign Central map**

$$\mathcal{F}_{V}^{(1)} = \sum_{i=1}^{V} \gamma_{i}^{(1)} x_{i} x_{(i \mod V)+1} \rightarrow \mathbf{F}_{V}^{(1)} = \begin{pmatrix} 0 & \gamma_{1}^{(1)} & 0 & \cdots & 0 \\ 0 & 0 & \gamma_{2}^{(1)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_{V-1}^{(1)} \\ \gamma_{V}^{(1)} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

$$\mathcal{F}_{V}^{(2)} = \sum_{i=1}^{V} \gamma_{i}^{(2)} x_{i} x_{(i+1 \mod V)+1} \rightarrow \mathbf{F}_{V}^{(2)} = \begin{pmatrix} 0 & 0 & \gamma_{1}^{(2)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_{V-1}^{(2)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_{V-2}^{(2)} \\ \gamma_{V-1}^{(2)} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

## **MQ-Sign Central map**

$$\mathcal{F}_{V}^{(1)} = \sum_{i=1}^{v} \gamma_{i}^{(1)} x_{i} x_{(i \mod v)+1} \rightarrow \mathbf{F}_{V}^{(1)} = \begin{pmatrix} 0 & \gamma_{1}^{(1)} & 0 & \cdots & 0 \\ 0 & 0 & \gamma_{2}^{(1)} & \cdots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & & \gamma_{v-1}^{(1)} \\ \gamma_{v}^{(1)} & 0 & 0 & \cdots & & 0 \end{pmatrix}$$

$$\mathcal{F}_{V}^{(2)} = \sum_{i=1}^{v} \gamma_{i}^{(2)} x_{i} x_{(i+1 \mod v)+1} \rightarrow \mathbf{F}_{V}^{(2)} = \begin{pmatrix} 0 & 0 & \gamma_{1}^{(2)} & \cdots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & & \gamma_{v-2}^{(2)} \\ \gamma_{v-1}^{(2)} & 0 & 0 & \cdots & & 0 \\ 0 & \gamma_{v}^{(2)} & 0 & \cdots & & 0 \end{pmatrix}$$

# **MQ-Sign Central map**

$$\mathcal{F}_{V}^{(1)} = \sum_{i=1}^{v} \gamma_{i}^{(1)} x_{i} x_{(i \bmod v)+1} \rightarrow \mathbf{F}_{V}^{(1)} = \begin{pmatrix} 0 & \gamma_{1}^{(1)} & 0 & \cdots & 0 \\ 0 & 0 & \gamma_{2}^{(1)} & \cdots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_{v-1}^{(1)} \\ \gamma_{v}^{(1)} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

$$\mathcal{F}_{V}^{(2)} = \sum_{i=1}^{v} \gamma_{i}^{(2)} x_{i} x_{i} x_{(i+1 \bmod v)+1} \rightarrow \mathbf{F}_{V}^{(2)} = \begin{pmatrix} 0 & 0 & \gamma_{1}^{(2)} & \cdots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_{v-2}^{(2)} \\ \gamma_{v-1}^{(2)} & 0 & 0 & \cdots & 0 \\ 0 & \gamma_{v}^{(2)} & 0 & \cdots & 0 \end{pmatrix}$$

21 / 23

Recall that the key equation  $\mathcal{P} = \mathcal{F} \circ \mathbf{S}$  translates to the matrix equations  $\mathbf{P^{(k)}} = \mathbf{S}^{\top} \mathbf{F^{(k)}} \mathbf{S}$ , i.e.

$$\begin{pmatrix} \mathbf{P}_1^{(k)} & \mathbf{P}_2^{(k)} \\ \mathbf{0} & \mathbf{P}_4^{(k)} \end{pmatrix} = \mathtt{Upper} \begin{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{S}_1^\top & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{F}_1^{(k)} & \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{S}_1 \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbf{F}_1^{(k)} & (\mathbf{F}_1^{(k)} + \mathbf{F}_1^{(k)\top}) \mathbf{S}_1 + \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathtt{Upper} & (\mathbf{S}_1^\top \mathbf{F}_1^{(k)} \mathbf{S}_1 + \mathbf{S}_1^\top \mathbf{F}_2^{(k)}) \end{pmatrix}.$$

$$egin{array}{lll} {f P}_1^{(k)} &=& {f F}_1^{(k)} & {
m and} \ {f P}_2^{(k)} &=& ({f P}_1^{(k)} + {f P}_1^{(k) op}) {f S}_1 + {f F}_2^{(k)}. \end{array}$$

- The second is a system of linear equations in the entries of the secret S<sub>1</sub>
- Still, not possible to determine them, due to the secret coefficients in  $\mathbf{F}_2^{(k)}$

Recall that the key equation  $\mathcal{P} = \mathcal{F} \circ \mathbf{S}$  translates to the matrix equations  $\mathbf{P}^{(k)} = \mathbf{S}^{\top} \mathbf{F}^{(k)} \mathbf{S}$ , i.e.

$$\begin{pmatrix} \mathbf{P}_1^{(k)} & \mathbf{P}_2^{(k)} \\ \mathbf{0} & \mathbf{P}_4^{(k)} \end{pmatrix} = \mathtt{Upper} \begin{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{S}_1^\top & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{F}_1^{(k)} & \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{S}_1 \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbf{F}_1^{(k)} & (\mathbf{F}_1^{(k)} + \mathbf{F}_1^{(k)\top})\mathbf{S}_1 + \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathtt{Upper} & (\mathbf{S}_1^\top \mathbf{F}_1^{(k)}\mathbf{S}_1 + \mathbf{S}_1^\top \mathbf{F}_2^{(k)}) \end{pmatrix}.$$

$$egin{array}{lll} {\sf P}_1^{(k)} &=& {\sf F}_1^{(k)} & {\sf and} \ {\sf P}_2^{(k)} &=& ({\sf P}_1^{(k)} + {\sf P}_1^{(k) op}) {\sf S}_1 + {\sf F}_2^{(k)}. \end{array}$$

- The second is a system of linear equations in the entries of the secret S<sub>1</sub>
- Still, not possible to determine them, due to the secret coefficients in  $\mathbf{F}_2^{(k)}$

Recall that the key equation  $\mathcal{P} = \mathcal{F} \circ \mathbf{S}$  translates to the matrix equations  $\mathbf{P}^{(k)} = \mathbf{S}^{\top} \mathbf{F}^{(k)} \mathbf{S}$ , i.e.

$$\begin{pmatrix} \mathbf{P}_1^{(k)} & \mathbf{P}_2^{(k)} \\ \mathbf{0} & \mathbf{P}_4^{(k)} \end{pmatrix} = \text{Upper} \begin{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{S}_1^\top & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{F}_1^{(k)} & \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{S}_1 \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbf{F}_1^{(k)} & (\mathbf{F}_1^{(k)} + \mathbf{F}_1^{(k)\top})\mathbf{S}_1 + \mathbf{F}_2^{(k)} \\ \mathbf{0} & \text{Upper} & (\mathbf{S}_1^\top \mathbf{F}_1^{(k)}\mathbf{S}_1 + \mathbf{S}_1^\top \mathbf{F}_2^{(k)}) \end{pmatrix}.$$

$$\begin{array}{lll} \textbf{P}_{1}^{(k)} & = & \textbf{F}_{1}^{(k)} & \text{and} \\ \textbf{P}_{2}^{(k)} & = & (\textbf{P}_{1}^{(k)} + \textbf{P}_{1}^{(k)\top}) \textbf{S}_{1} + \textbf{F}_{2}^{(k)}. \end{array}$$

- The second is a system of linear equations in the entries of the secret S<sub>1</sub>
- Still, not possible to determine them, due to the secret coefficients in  $\mathbf{F}_2^{(k)}$

Recall that the key equation  $\mathcal{P} = \mathcal{F} \circ \mathbf{S}$  translates to the matrix equations  $\mathbf{P}^{(k)} = \mathbf{S}^{\top} \mathbf{F}^{(k)} \mathbf{S}$ , i.e.

$$\begin{pmatrix} \mathbf{P}_1^{(k)} & \mathbf{P}_2^{(k)} \\ \mathbf{0} & \mathbf{P}_4^{(k)} \end{pmatrix} = \text{Upper} \begin{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{S}_1^\top & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{F}_1^{(k)} & \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{S}_1 \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbf{F}_1^{(k)} & (\mathbf{F}_1^{(k)} + \mathbf{F}_1^{(k)\top})\mathbf{S}_1 + \mathbf{F}_2^{(k)} \\ \mathbf{0} & \text{Upper} & (\mathbf{S}_1^\top \mathbf{F}_1^{(k)}\mathbf{S}_1 + \mathbf{S}_1^\top \mathbf{F}_2^{(k)}) \end{pmatrix}.$$

$$\begin{array}{lcl} \textbf{P}_{1}^{(k)} & = & \textbf{F}_{1}^{(k)} & \text{and} \\ \textbf{P}_{2}^{(k)} & = & (\textbf{P}_{1}^{(k)} + \textbf{P}_{1}^{(k)\top}) \textbf{S}_{1} + \textbf{F}_{2}^{(k)}. \end{array}$$

- The second is a system of linear equations in the entries of the secret  $S_1$
- Still, not possible to determine them, due to the secret coefficients in  $\mathbf{F}_2^{(k)}$

Recall that the key equation  $\mathcal{P} = \mathcal{F} \circ \mathbf{S}$  translates to the matrix equations  $\mathbf{P}^{(k)} = \mathbf{S}^{\top} \mathbf{F}^{(k)} \mathbf{S}$ , i.e.

$$\begin{pmatrix} \mathbf{P}_1^{(k)} & \mathbf{P}_2^{(k)} \\ \mathbf{0} & \mathbf{P}_4^{(k)} \end{pmatrix} = \text{Upper} \begin{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{S}_1^\top & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{F}_1^{(k)} & \mathbf{F}_2^{(k)} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{I} & \mathbf{S}_1 \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} \mathbf{F}_1^{(k)} & (\mathbf{F}_1^{(k)} + \mathbf{F}_1^{(k)\top})\mathbf{S}_1 + \mathbf{F}_2^{(k)} \\ \mathbf{0} & \text{Upper} & (\mathbf{S}_1^\top \mathbf{F}_1^{(k)}\mathbf{S}_1 + \mathbf{S}_1^\top \mathbf{F}_2^{(k)}) \end{pmatrix}.$$

$$\begin{array}{lll} \textbf{P}_{1}^{(k)} & = & \textbf{F}_{1}^{(k)} & \text{and} \\ \textbf{P}_{2}^{(k)} & = & (\textbf{P}_{1}^{(k)} + \textbf{P}_{1}^{(k)\top}) \textbf{S}_{1} + \textbf{F}_{2}^{(k)}. \end{array}$$

- The second is a system of linear equations in the entries of the secret S<sub>1</sub>
- Still, not possible to determine them, due to the secret coefficients in  $\mathbf{F}_2^{(k)}$

In two variants of MQ-Sign, the coefficients in  $F_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\mathbf{P}_{2}^{(k)} = (\mathbf{P}_{1}^{(k)} + \mathbf{P}_{1}^{(k)\top})\mathbf{S}_{1} + \mathbf{F}_{2}^{(k)}.$$

$$\underbrace{\begin{pmatrix} p_{1,v+1}^{(k)} & \cdots & p_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ p_{v,v+1}^{(k)} & \cdots & p_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{\begin{pmatrix} p_{1,1}^{(k)} & \cdots & p_{1,v}^{(k)} \\ \vdots & & \vdots \\ p_{v,1}^{(k)} & \cdots & p_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{\begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix}}_{secret} + \underbrace{\begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots &$$

secret, but known structure

In two variants of MQ-Sign, the coefficients in  $F_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\mathbf{P}_{2}^{(k)} = (\mathbf{P}_{1}^{(k)} + \mathbf{P}_{1}^{(k)\top})\mathbf{S}_{1} + \mathbf{F}_{2}^{(k)}.$$

$$\underbrace{\begin{pmatrix} \boldsymbol{p}_{1,v+1}^{(k)} & \cdots & \boldsymbol{p}_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ \boldsymbol{p}_{v,v+1}^{(k)} & \cdots & \boldsymbol{p}_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{\begin{pmatrix} \boldsymbol{p}_{1,1}^{(k)} & \cdots & \boldsymbol{p}_{1,v}^{(k)} \\ \vdots & & \vdots \\ \boldsymbol{p}_{v,1}^{(k)} & \cdots & \boldsymbol{p}_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{\begin{pmatrix} \boldsymbol{s}_{11} & \cdots & \boldsymbol{s}_{1m} \\ \vdots & & \vdots \\ \boldsymbol{s}_{v1} & \cdots & \boldsymbol{s}_{vm} \end{pmatrix}}_{secret} + \underbrace{\begin{pmatrix} \boldsymbol{0} & \boldsymbol{\gamma}_{1}^{(k)} & \cdots & \boldsymbol{0} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{0} & \boldsymbol{0} & \cdots & \boldsymbol{\gamma}_{m-1}^{(k)} \\ \boldsymbol{\gamma}_{m}^{(k)} & \boldsymbol{0} & \cdots & \boldsymbol{0} \\ \vdots & \vdots & \ddots & \cdots & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{0} & \cdots & \boldsymbol{0} \end{pmatrix}}_{secret, but known structure}$$

In two variants of MQ-Sign, the coefficients in  $\mathbf{F}_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\mathbf{P}_2^{(k)} = (\mathbf{P}_1^{(k)} + \mathbf{P}_1^{(k)\top})\mathbf{S}_1 + \mathbf{F}_2^{(k)}.$$

$$\underbrace{\begin{pmatrix} p_{1,v+1}^{(k)} & \cdots & p_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ p_{v,v+1}^{(k)} & \cdots & p_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{\begin{pmatrix} p_{1,1}^{(k)} & \cdots & p_{1,v}^{(k)} \\ \vdots & & \vdots \\ p_{v,1}^{(k)} & \cdots & p_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{\begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix}}_{secret} + \underbrace{\begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \vdots & \vdots & \vdots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ \vdots$$

secret, but known structure

In two variants of MQ-Sign, the coefficients in  $F_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\mathbf{P}_{2}^{(k)} = (\mathbf{P}_{1}^{(k)} + \mathbf{P}_{1}^{(k)\top})\mathbf{S}_{1} + \mathbf{F}_{2}^{(k)}.$$

$$\underbrace{\begin{pmatrix} \rho_{1,v+1}^{(k)} & \cdots & \rho_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ \rho_{v,v+1}^{(k)} & \cdots & \rho_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{\begin{pmatrix} \rho_{1,1}^{(k)} & \cdots & \rho_{1,v}^{(k)} \\ \vdots & & \vdots \\ \rho_{v,1}^{(k)} & \cdots & \rho_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{\begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix}}_{secret} + \underbrace{\begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \gamma_{m}^{(k)} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots &$$

In two variants of MQ-Sign, the coefficients in  $\mathbf{F}_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\underbrace{ \begin{pmatrix} p_{1,v+1}^{(k)} & \cdots & p_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ p_{v,v+1}^{(k)} & \cdots & p_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{ \begin{pmatrix} p_{1,1}^{(k)} & \cdots & p_{1,v}^{(k)} \\ \vdots & & \vdots \\ p_{v,1}^{(k)} & \cdots & p_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{ \begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix}}_{secret} + \underbrace{ \begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots$$

- Collect linear equations for all  $k \in \{1, \dots, m\}$  polynomials.
- Obtain system of mv(m-1) equations in vm variables (can be divided into subsystems).
- Once S is known, the central polynomials can efficiently be found

The constructed key is actually not equivalent to a UOV key, it is weaker!

In two variants of MQ-Sign, the coefficients in  $\mathbf{F}_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\underbrace{ \begin{pmatrix} p_{1,v+1}^{(k)} & \cdots & p_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ p_{v,v+1}^{(k)} & \cdots & p_{v,v+m}^{(k)} \end{pmatrix}}_{public} = \underbrace{ \begin{pmatrix} p_{1,1}^{(k)} & \cdots & p_{1,v}^{(k)} \\ \vdots & & \vdots \\ p_{v,1}^{(k)} & \cdots & p_{v,v}^{(k)} \end{pmatrix}}_{public} \underbrace{ \begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix}}_{secret} + \underbrace{ \begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \vdots & \vdots & \vdots \\ 0 & \vdots & \vdots & \vdots \\ 0 & \vdots$$

- Collect linear equations for all  $k \in \{1, ..., m\}$  polynomials.
- Obtain system of mv(m-1) equations in vm variables (can be divided into subsystems).
- Once **S** is known, the central polynomials can efficiently be found.

The constructed key is actually not equivalent to a UOV key, it is weaker!

In two variants of MQ-Sign, the coefficients in  $F_2^{(k)}$  are chosen sparsely.

This removes unknown variables from the system

$$\underbrace{ \begin{pmatrix} p_{1,v+1}^{(k)} & \cdots & p_{1,v+m}^{(k)} \\ \vdots & & \vdots \\ p_{v,v+1}^{(k)} & \cdots & p_{v,v+m}^{(k)} \end{pmatrix} }_{public} = \underbrace{ \begin{pmatrix} p_{1,1}^{(k)} & \cdots & p_{1,v}^{(k)} \\ \vdots & & \vdots \\ p_{v,1}^{(k)} & \cdots & p_{v,v}^{(k)} \end{pmatrix} }_{public} \underbrace{ \begin{pmatrix} s_{11} & \cdots & s_{1m} \\ \vdots & & \vdots \\ s_{v1} & \cdots & s_{vm} \end{pmatrix} }_{secret} + \underbrace{ \begin{pmatrix} 0 & \gamma_{1}^{(k)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \gamma_{m-1}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots$$

- Collect linear equations for all  $k \in \{1, ..., m\}$  polynomials.
- Obtain system of mv(m-1) equations in vm variables (can be divided into subsystems).
- Once **S** is known, the central polynomials can efficiently be found.

The constructed key is actually not equivalent to a UOV key, it is weaker!